

Measuring Inconsistency in Knowledge Bases

John Grant and Anthony Hunter

George Voutsadakis¹

¹Mathematics and Computer Science
Lake Superior State University

Seminar Presentation

Universidade do Minho/Universidade de Aveiro

1 Measuring Inconsistency In Knowledge Bases

- Introduction
- First-Order QC Logic: The Syntax
- First-Order QC Logic: The Semantics
- QC Models
- The Inconsistency Measure
- Extrinsic Inconsistency

Main Points

Main Points

- Knowledge Bases often contain inconsistencies;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;
 - first-order statements representing knowledge (or database) integrity constraints imposed on the data;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;
 - first-order statements representing knowledge (or database) integrity constraints imposed on the data;
- To measure the “extrinsic” inconsistency a comparison is made across different knowledge bases;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;
 - first-order statements representing knowledge (or database) integrity constraints imposed on the data;
- To measure the “extrinsic” inconsistency a comparison is made across different knowledge bases;
- The measure depends on

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;
 - first-order statements representing knowledge (or database) integrity constraints imposed on the data;
- To measure the “extrinsic” inconsistency a comparison is made across different knowledge bases;
- The measure depends on
 - the language used (relation symbols and constants);

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;
 - first-order statements representing knowledge (or database) integrity constraints imposed on the data;
- To measure the “extrinsic” inconsistency a comparison is made across different knowledge bases;
- The measure depends on
 - the language used (relation symbols and constants);
 - the underlying domain of individuals;

Main Points

- Knowledge Bases often contain inconsistencies;
- We want to have a measure for comparing the inconsistencies arising in various knowledge bases;
- The platform used is Quasi-classical (QC) logic;
- There are
 - constants representing individuals in the knowledge base;
 - first-order statements representing knowledge (or database) integrity constraints imposed on the data;
- To measure the “extrinsic” inconsistency a comparison is made across different knowledge bases;
- The measure depends on
 - the language used (relation symbols and constants);
 - the underlying domain of individuals;
- It is shown that this measure accomplishes also an evaluation of the “intrinsic” inconsistency of a knowledge base;

Introducing Quasi-Classical (QC) Logic

Introducing Quasi-Classical (QC) Logic

- Vocabulary:

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;
 - Predicate Symbols \mathcal{P} ;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;
 - Predicate Symbols \mathcal{P} ;
 - We assume that **no non-nullary functions symbols** are available;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;
 - Predicate Symbols \mathcal{P} ;
 - We assume that **no non-nullary functions symbols** are available;
- We use

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;
 - Predicate Symbols \mathcal{P} ;
 - We assume that **no non-nullary functions symbols** are available;
- We use
 - P, R, \dots for predicate symbols;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;
 - Predicate Symbols \mathcal{P} ;
 - We assume that **no non-nullary functions symbols** are available;
- We use
 - P, R, \dots for predicate symbols;
 - a, b, c, \dots for constant symbols;

Introducing Quasi-Classical (QC) Logic

- Vocabulary:
 - The logical symbols:
 - A countable set of variable symbols \mathcal{V} ;
 - The ordinary connectives $\{\neg, \vee, \wedge, \rightarrow\}$;
 - The two quantifiers $\{\forall, \exists\}$;
 - Punctuation symbols for readability;
 - The extra-logical symbols:
 - Constant symbols \mathcal{C} ;
 - Predicate Symbols \mathcal{P} ;
 - We assume that **no non-nullary functions symbols** are available;
- We use
 - P, R, \dots for predicate symbols;
 - a, b, c, \dots for constant symbols;
 - x, y, \dots for individual variables;

Syntax

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;
- Usage is made of

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;
- Usage is made of
 - $\alpha, \beta, \gamma, \dots$ for literals;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;
- Usage is made of
 - $\alpha, \beta, \gamma, \dots$ for literals;
 - ϕ for a **clause** (disjunction of literals);

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;
- Usage is made of
 - $\alpha, \beta, \gamma, \dots$ for literals;
 - ϕ for a **clause** (disjunction of literals);
 - ψ for conjunction of clauses;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;
- Usage is made of
 - $\alpha, \beta, \gamma, \dots$ for literals;
 - ϕ for a **clause** (disjunction of literals);
 - ψ for conjunction of clauses;
 - θ for arbitrary formulas;

Syntax

- As a consequence of the language choices a **term** is either a variable or a constant symbol;
- The following metalogical notions are defined as usual:
 - **free variable**,
 - **bound variable**,
 - **ground formula**;
- An **atom** is of the form $P(t_1, \dots, t_n)$, where t_1, \dots, t_n are terms;
- A **literal** is an atom or the negation of an atom;
- **Formulas** are defined by the usual recursive rules starting with atomic formulas and using the connectives and the quantifiers;
- Usage is made of
 - $\alpha, \beta, \gamma, \dots$ for literals;
 - ϕ for a **clause** (disjunction of literals);
 - ψ for conjunction of clauses;
 - θ for arbitrary formulas;
- The **set of \mathcal{L} -formulas** is denoted by $\text{Fm}(\mathcal{L})$;

Complementation and Focus

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;
- Let $\alpha_j, j = 1, \dots, n$ be literals;

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;
- Let $\alpha_j, j = 1, \dots, n$ be literals; Consider the clause $\alpha_1 \vee \dots \vee \alpha_n$;

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;
- Let $\alpha_i, i = 1, \dots, n$ be literals; Consider the clause $\alpha_1 \vee \dots \vee \alpha_n$; The **focus of** $\alpha_1 \vee \dots \vee \alpha_n$ **by** α_i is defined by

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;
- Let $\alpha_i, i = 1, \dots, n$ be literals; Consider the clause $\alpha_1 \vee \dots \vee \alpha_n$; The **focus of** $\alpha_1 \vee \dots \vee \alpha_n$ **by** α_i is defined by

$$\otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i) := \alpha_1 \vee \dots \vee \alpha_{i-1} \vee \alpha_{i+1} \vee \dots \vee \alpha_n;$$

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;
- Let $\alpha_i, i = 1, \dots, n$ be literals; Consider the clause $\alpha_1 \vee \dots \vee \alpha_n$; The **focus of** $\alpha_1 \vee \dots \vee \alpha_n$ **by** α_i is defined by

$$\otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i) := \alpha_1 \vee \dots \vee \alpha_{i-1} \vee \alpha_{i+1} \vee \dots \vee \alpha_n;$$

- Example:** If α, β, γ are literals, then

$$\otimes(\alpha \vee \beta \vee \gamma, \beta) =$$

Complementation and Focus

- Given an atom α one obtains two associated literals α and $\neg\alpha$;
- The **complementation** \sim applied to a literal is defined by

$$\sim \alpha := \neg\alpha \quad \text{and} \quad \sim (\neg\alpha) := \alpha;$$

- Note that complementation is a **metalinguistic** convenience;
- Let $\alpha_i, i = 1, \dots, n$ be literals; Consider the clause $\alpha_1 \vee \dots \vee \alpha_n$; The **focus of** $\alpha_1 \vee \dots \vee \alpha_n$ **by** α_i is defined by

$$\otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i) := \alpha_1 \vee \dots \vee \alpha_{i-1} \vee \alpha_{i+1} \vee \dots \vee \alpha_n;$$

- Example:** If α, β, γ are literals, then

$$\otimes(\alpha \vee \beta \vee \gamma, \beta) = \alpha \vee \gamma;$$

\mathcal{L} -Structures

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:
 - For $c \in \mathcal{C}$, $I(c) \in D$;

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:
 - For $c \in \mathcal{C}$, $I(c) \in D$;
 - For $P(n) \in \mathcal{P}$ (an n -ary predicate symbol) $I(P) : D^n \rightarrow \{0, 1\}$ is an n -ary predicate over D ;

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:
 - For $c \in \mathcal{C}$, $I(c) \in D$;
 - For $P(n) \in \mathcal{P}$ (an n -ary predicate symbol) $I(P) : D^n \rightarrow \{0, 1\}$ is an n -ary predicate over D ;
- The following **important convention** will be followed to simplify the treatment:

I maps constant symbols to themselves.

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:
 - For $c \in \mathcal{C}$, $I(c) \in D$;
 - For $P(n) \in \mathcal{P}$ (an n -ary predicate symbol) $I(P) : D^n \rightarrow \{0, 1\}$ is an n -ary predicate over D ;
- The following **important convention** will be followed to simplify the treatment:

I maps constant symbols to themselves.
- This assumption forces the consideration of languages $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$, such that $\mathcal{C} \subseteq D$, where D is the domain of interpretation of the \mathcal{L} -structures considered;

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:
 - For $c \in \mathcal{C}$, $I(c) \in D$;
 - For $P(n) \in \mathcal{P}$ (an n -ary predicate symbol) $I(P) : D^n \rightarrow \{0, 1\}$ is an n -ary predicate over D ;
- The following **important convention** will be followed to simplify the treatment:

I maps constant symbols to themselves.

- This assumption forces the consideration of languages $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$, such that $\mathcal{C} \subseteq D$, where D is the domain of interpretation of the \mathcal{L} -structures considered;
- Going even further, by introducing a new constant for each element in $D \setminus \mathcal{C}$, we assume that $\mathcal{C} = D$;

\mathcal{L} -Structures

- A **classical structure** for a language \mathcal{L} is a pair (D, I) , where
 - $D \neq \emptyset$ is the **domain**;
 - I is the **interpretation** function, assigning elements and predicates over D to the symbols in \mathcal{L} as follows:
 - For $c \in \mathcal{C}$, $I(c) \in D$;
 - For $P(n) \in \mathcal{P}$ (an n -ary predicate symbol) $I(P) : D^n \rightarrow \{0, 1\}$ is an n -ary predicate over D ;
- The following **important convention** will be followed to simplify the treatment:

I maps constant symbols to themselves.

- This assumption forces the consideration of languages $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$, such that $\mathcal{C} \subseteq D$, where D is the domain of interpretation of the \mathcal{L} -structures considered;
- Going even further, by introducing a new constant for each element in $D \setminus \mathcal{C}$, we assume that $\mathcal{C} = D$; Thus, we work with languages $\mathcal{L} = \langle D, \mathcal{P} \rangle$, where D is the domain of interpretation;

Assignments and Substitutions

Assignments and Substitutions

- Let $\mathcal{L} = \langle D, \mathcal{P} \rangle$ be a language;

Assignments and Substitutions

- Let $\mathcal{L} = \langle D, \mathcal{P} \rangle$ be a language;
- Let (D, I) be an \mathcal{L} -structure;

Assignments and Substitutions

- Let $\mathcal{L} = \langle D, \mathcal{P} \rangle$ be a language;
- Let (D, I) be an \mathcal{L} -structure;
- An **assignment** is a function $A : \mathcal{V} \rightarrow D$;

Assignments and Substitutions

- Let $\mathcal{L} = \langle D, \mathcal{P} \rangle$ be a language;
- Let (D, I) be an \mathcal{L} -structure;
- An **assignment** is a function $A : \mathcal{V} \rightarrow D$;
- Given an assignment A and $x \in \mathcal{V}$, an **x -variant assignment** A' is an assignment such that

$$A'(y) = A(y), \text{ for all } y \in \mathcal{V}, y \neq x;$$

Assignments and Substitutions

- Let $\mathcal{L} = \langle D, \mathcal{P} \rangle$ be a language;
- Let (D, I) be an \mathcal{L} -structure;
- An **assignment** is a function $A : \mathcal{V} \rightarrow D$;
- Given an assignment A and $x \in \mathcal{V}$, an **x -variant assignment** A' is an assignment such that

$$A'(y) = A(y), \text{ for all } y \in \mathcal{V}, y \neq x;$$

- The **extension** A (abusing notation) of an assignment to terms is defined by

$$A(t) = \begin{cases} d, & \text{if } t = x \text{ and } A(x) = d \\ d, & \text{if } t = d \end{cases}$$

Assignments and Substitutions

- Let $\mathcal{L} = \langle D, \mathcal{P} \rangle$ be a language;
- Let (D, I) be an \mathcal{L} -structure;
- An **assignment** is a function $A : \mathcal{V} \rightarrow D$;
- Given an assignment A and $x \in \mathcal{V}$, an **x -variant assignment** A' is an assignment such that

$$A'(y) = A(y), \text{ for all } y \in \mathcal{V}, y \neq x;$$

- The **extension** A (abusing notation) of an assignment to terms is defined by

$$A(t) = \begin{cases} d, & \text{if } t = x \text{ and } A(x) = d \\ d, & \text{if } t = d \end{cases}$$

- For a formula θ in which $x \in \mathcal{V}$ occurs free, and a constant $d \in D$, define the **substitution** $\theta[x/d]$ of x by d in θ as the formula obtained by replacing all occurrences of the free variable x by d ;

Quasi-Classical Bistructures

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;
- Since $I^+(c) = I^-(c) = c$, I^+ and I^- can only differ in the interpretations of the predicate symbols;

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;
- Since $I^+(c) = I^-(c) = c$, I^+ and I^- can only differ in the interpretations of the predicate symbols;
- Intuitively, we conceive of I^+ as the interpretation of positive literals and of I^- as the interpretation of negative literals;

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;
- Since $I^+(c) = I^-(c) = c$, I^+ and I^- can only differ in the interpretations of the predicate symbols;
- Intuitively, we conceive of I^+ as the interpretation of positive literals and of I^- as the interpretation of negative literals; This intuition is formalized as follows:

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;
- Since $I^+(c) = I^-(c) = c$, I^+ and I^- can only differ in the interpretations of the predicate symbols;
- Intuitively, we conceive of I^+ as the interpretation of positive literals and of I^- as the interpretation of negative literals; This intuition is formalized as follows:
- Given a bistructure $E = (D, I^+, I^-)$ and an assignment A , the **decoupled satisfaction** \models_d for literals in $\mathcal{L}(D)$ is defined as follows:

$$(E, A) \models_d \alpha \quad \text{iff} \quad \begin{cases} \alpha = P(t_1, \dots, t_n) \text{ and } I^+(P)(A(t_1), \dots, A(t_n)) = 1 \text{ or} \\ \alpha = \neg P(t_1, \dots, t_n) \text{ and } I^-(P)(A(t_1), \dots, A(t_n)) = 1 \end{cases}$$

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;
- Since $I^+(c) = I^-(c) = c$, I^+ and I^- can only differ in the interpretations of the predicate symbols;
- Intuitively, we conceive of I^+ as the interpretation of positive literals and of I^- as the interpretation of negative literals; This intuition is formalized as follows:
- Given a bistructure $E = (D, I^+, I^-)$ and an assignment A , the **decoupled satisfaction** \models_d for literals in $\mathcal{L}(D)$ is defined as follows:

$$(E, A) \models_d \alpha \quad \text{iff} \quad \begin{cases} \alpha = P(t_1, \dots, t_n) \text{ and } I^+(P)(A(t_1), \dots, A(t_n)) = 1 \text{ or} \\ \alpha = \neg P(t_1, \dots, t_n) \text{ and } I^-(P)(A(t_1), \dots, A(t_n)) = 1 \end{cases}$$

- Note that both an atom and its negation may be true in a bistructure;

Quasi-Classical Bistructures

- An \mathcal{L} -**bistructure** is a tuple (D, I^+, I^-) , where (D, I^+) and (D, I^-) are classical \mathcal{L} -structures;
- Since $I^+(c) = I^-(c) = c$, I^+ and I^- can only differ in the interpretations of the predicate symbols;
- Intuitively, we conceive of I^+ as the interpretation of positive literals and of I^- as the interpretation of negative literals; This intuition is formalized as follows:
- Given a bistructure $E = (D, I^+, I^-)$ and an assignment A , the **decoupled satisfaction** \models_d for literals in $\mathcal{L}(D)$ is defined as follows:

$$(E, A) \models_d \alpha \quad \text{iff} \quad \begin{cases} \alpha = P(t_1, \dots, t_n) \text{ and } I^+(P)(A(t_1), \dots, A(t_n)) = 1 \text{ or} \\ \alpha = \neg P(t_1, \dots, t_n) \text{ and } I^-(P)(A(t_1), \dots, A(t_n)) = 1 \end{cases}$$

- Note that both an atom and its negation may be true in a bistructure;
- This constitutes the justification for the term *decoupled satisfaction* and, also, provides the basis for **paraconsistent reasoning**;

Prenex Conjunctive Normal Form (PCNF) Hypothesis

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Classical PCNF Theorem

Every formula is logically equivalent using classical entailment to one in prenex conjunctive normal form.

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Classical PCNF Theorem

Every formula is logically equivalent using classical entailment to one in prenex conjunctive normal form.

- The process of converting a given formula to PCNF involves

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Classical PCNF Theorem

Every formula is logically equivalent using classical entailment to one in prenex conjunctive normal form.

- The process of converting a given formula to PCNF involves
 - moving quantifiers at the front;

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Classical PCNF Theorem

Every formula is logically equivalent using classical entailment to one in prenex conjunctive normal form.

- The process of converting a given formula to PCNF involves
 - moving quantifiers at the front;
 - converting the matrix into CNF using the commutative, associative and distributive laws;

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Classical PCNF Theorem

Every formula is logically equivalent using classical entailment to one in prenex conjunctive normal form.

- The process of converting a given formula to PCNF involves
 - moving quantifiers at the front;
 - converting the matrix into CNF using the commutative, associative and distributive laws;
 - rewriting \rightarrow using \neg and \vee ;

Prenex Conjunctive Normal Form (PCNF) Hypothesis

- In the sequel, the assumption is that all formulas are in the prenex conjunctive normal form (PCNF):
 - all quantifiers appear at the front;
 - the matrix consists of a conjunction of clauses or a single clause;

Classical PCNF Theorem

Every formula is logically equivalent using classical entailment to one in prenex conjunctive normal form.

- The process of converting a given formula to PCNF involves
 - moving quantifiers at the front;
 - converting the matrix into CNF using the commutative, associative and distributive laws;
 - rewriting \rightarrow using \neg and \vee ;
- In the present setting **no rules involving TAUTOLOGIES or CONTRADICTIONS are allowed** in the rewriting process!

Strong Satisfaction

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:
 - $(E, A) \models_s \alpha$ iff $(E, A) \models_d \alpha$, for α a literal;

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:
 - $(E, A) \models_s \alpha$ iff $(E, A) \models_d \alpha$, for α a literal;
 - $(E, A) \models_s \alpha_1 \vee \dots \vee \alpha_n$ iff

$[(E, A) \models_s \alpha_1 \text{ or } \dots \text{ or } (E, A) \models_s \alpha_n]$ and, for all $1 \leq i \leq n$,
 $[(E, A) \models_{s\sim} \alpha_i \text{ implies } (E, A) \models_s \otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i)]$;

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:

- $(E, A) \models_s \alpha$ iff $(E, A) \models_d \alpha$, for α a literal;
- $(E, A) \models_s \alpha_1 \vee \dots \vee \alpha_n$ iff

$[(E, A) \models_s \alpha_1 \text{ or } \dots \text{ or } (E, A) \models_s \alpha_n]$ and, for all $1 \leq i \leq n$,
 $[(E, A) \models_{s\sim} \alpha_i \text{ implies } (E, A) \models_s \otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i)]$;

- $(E, A) \models_s \phi_1 \wedge \dots \wedge \phi_m$ iff $[(E, A) \models_s \phi_1 \text{ and } \dots \text{ and } (E, A) \models_s \phi_m]$;

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:

- $(E, A) \models_s \alpha$ iff $(E, A) \models_d \alpha$, for α a literal;
- $(E, A) \models_s \alpha_1 \vee \dots \vee \alpha_n$ iff

$$[(E, A) \models_s \alpha_1 \text{ or } \dots \text{ or } (E, A) \models_s \alpha_n] \text{ and, for all } 1 \leq i \leq n, \\ [(E, A) \models_s \sim \alpha_i \text{ implies } (E, A) \models_s \otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i)];$$

- $(E, A) \models_s \phi_1 \wedge \dots \wedge \phi_m$ iff $[(E, A) \models_s \phi_1 \text{ and } \dots \text{ and } (E, A) \models_s \phi_m]$;
- $(E, A) \models_s \exists x \theta$ iff, for some x -variant A' , $(E, A') \models_s \theta[x/A'(x)]$;

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:

- $(E, A) \models_s \alpha$ iff $(E, A) \models_d \alpha$, for α a literal;
- $(E, A) \models_s \alpha_1 \vee \dots \vee \alpha_n$ iff

$[(E, A) \models_s \alpha_1 \text{ or } \dots \text{ or } (E, A) \models_s \alpha_n]$ and, for all $1 \leq i \leq n$,
 $[(E, A) \models_s \sim \alpha_i \text{ implies } (E, A) \models_s \otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i)]$;

- $(E, A) \models_s \phi_1 \wedge \dots \wedge \phi_m$ iff $[(E, A) \models_s \phi_1 \text{ and } \dots \text{ and } (E, A) \models_s \phi_m]$;
- $(E, A) \models_s \exists x\theta$ iff, for some x -variant A' , $(E, A') \models_s \theta[x/A'(x)]$;
- $(E, A) \models_s \forall x\theta$ iff, for all x -variant A' , $(E, A') \models_s \theta[x/A'(x)]$;

Strong Satisfaction

- The **strong satisfaction** \models_s is defined by structural induction on formulas, as follows:

- $(E, A) \models_s \alpha$ iff $(E, A) \models_d \alpha$, for α a literal;
- $(E, A) \models_s \alpha_1 \vee \dots \vee \alpha_n$ iff

$[(E, A) \models_s \alpha_1 \text{ or } \dots \text{ or } (E, A) \models_s \alpha_n]$ and, for all $1 \leq i \leq n$,
 $[(E, A) \models_s \sim \alpha_i \text{ implies } (E, A) \models_s \otimes(\alpha_1 \vee \dots \vee \alpha_n, \alpha_i)]$;

- $(E, A) \models_s \phi_1 \wedge \dots \wedge \phi_m$ iff $[(E, A) \models_s \phi_1 \text{ and } \dots \text{ and } (E, A) \models_s \phi_m]$;
- $(E, A) \models_s \exists x \theta$ iff, for some x -variant A' , $(E, A') \models_s \theta[x/A'(x)]$;
- $(E, A) \models_s \forall x \theta$ iff, for all x -variant A' , $(E, A') \models_s \theta[x/A'(x)]$;
- We write $E \models_s \theta$ to mean that $(E, A) \models_s \theta$, for all assignments A ;

QC Models

QC Models

- Let Δ be a finite set of formulae and E a bistructure;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;
 - $\text{GrdLt}(\mathcal{L}, D) = \text{GrdAt}(\mathcal{L}, D) \cup \{\neg\alpha : \alpha \in \text{GrdAt}(\mathcal{L}, D)\}$;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;
 - $\text{GrdLt}(\mathcal{L}, D) = \text{GrdAt}(\mathcal{L}, D) \cup \{\neg\alpha : \alpha \in \text{GrdAt}(\mathcal{L}, D)\}$;
- It is fairly obvious that a QC model $E = (D, I^+, I^-)$ may be equivalently represented by a set of ground literals as follows:

$$\{\alpha : E \models_s \alpha \text{ and } \alpha \in \text{GrdLt}(\mathcal{L}, D)\};$$

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;
 - $\text{GrdLt}(\mathcal{L}, D) = \text{GrdAt}(\mathcal{L}, D) \cup \{\neg\alpha : \alpha \in \text{GrdAt}(\mathcal{L}, D)\}$;
- It is fairly obvious that a QC model $E = (D, I^+, I^-)$ may be equivalently represented by a set of ground literals as follows:

$$\{\alpha : E \models_s \alpha \text{ and } \alpha \in \text{GrdLt}(\mathcal{L}, D)\};$$

- The following notation will be used:

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;
 - $\text{GrdLt}(\mathcal{L}, D) = \text{GrdAt}(\mathcal{L}, D) \cup \{\neg\alpha : \alpha \in \text{GrdAt}(\mathcal{L}, D)\}$;
- It is fairly obvious that a QC model $E = (D, I^+, I^-)$ may be equivalently represented by a set of ground literals as follows:

$$\{\alpha : E \models_s \alpha \text{ and } \alpha \in \text{GrdLt}(\mathcal{L}, D)\};$$

- The following notation will be used:
 - $\text{QC}(\mathcal{L}, \Delta, D)$ is the class of QC models of Δ , where the formulas are in \mathcal{L} and the domain of the QC models is D ;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;
 - $\text{GrdLt}(\mathcal{L}, D) = \text{GrdAt}(\mathcal{L}, D) \cup \{\neg\alpha : \alpha \in \text{GrdAt}(\mathcal{L}, D)\}$;
- It is fairly obvious that a QC model $E = (D, I^+, I^-)$ may be equivalently represented by a set of ground literals as follows:

$$\{\alpha : E \models_s \alpha \text{ and } \alpha \in \text{GrdLt}(\mathcal{L}, D)\};$$

- The following notation will be used:
 - $\text{QC}(\mathcal{L}, \Delta, D)$ is the class of QC models of Δ , where the formulas are in \mathcal{L} and the domain of the QC models is D ;
 - M denotes a QC model in the form of ground literals;

QC Models

- Let Δ be a finite set of formulae and E a bistructure; E is a **QC model** of Δ iff, for all $\theta \in \Delta$, $E \models_s \theta$;
- Let $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ be a language and D a domain; We define
 - $\text{GrdAt}(\mathcal{L}, D) = \{P(d_1, \dots, d_n) : P(n) \in \mathcal{P} \text{ and } d_1, \dots, d_n \in D\}$;
 - $\text{GrdLt}(\mathcal{L}, D) = \text{GrdAt}(\mathcal{L}, D) \cup \{\neg\alpha : \alpha \in \text{GrdAt}(\mathcal{L}, D)\}$;
- It is fairly obvious that a QC model $E = (D, I^+, I^-)$ may be equivalently represented by a set of ground literals as follows:

$$\{\alpha : E \models_s \alpha \text{ and } \alpha \in \text{GrdLt}(\mathcal{L}, D)\};$$

- The following notation will be used:
 - $\text{QC}(\mathcal{L}, \Delta, D)$ is the class of QC models of Δ , where the formulas are in \mathcal{L} and the domain of the QC models is D ;
 - M denotes a QC model in the form of ground literals; Of course this expressibility rests on the hypothesis $\mathcal{C} \subseteq D$;

Set of Satisfied Formulas

Set of Satisfied Formulas

- In the sequel **all languages $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ and all domains D are assumed finite;**

Set of Satisfied Formulas

- In the sequel **all languages $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ and all domains D are assumed finite**;
- Let \mathcal{L} be a language, $\{M_1, \dots, M_m\}$ a set of QC models, such that, for all i , $M_i \subseteq \text{GrdLt}(\mathcal{L}, D)$;

Set of Satisfied Formulas

- In the sequel **all languages** $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ and all domains D are assumed **finite**;
- Let \mathcal{L} be a language, $\{M_1, \dots, M_m\}$ a set of QC models, such that, for all i , $M_i \subseteq \text{GrdLt}(\mathcal{L}, D)$; The set of all **satisfied formulas** is

$$\text{SF}(\{M_1, \dots, M_n\}) := \{\theta \in \text{Fm}(\mathcal{L}) : \text{for all } i, M_i \models_s \theta\};$$

Set of Satisfied Formulas

- In the sequel **all languages** $\mathcal{L} = \langle \mathcal{C}, \mathcal{P} \rangle$ and all domains D are assumed **finite**;
- Let \mathcal{L} be a language, $\{M_1, \dots, M_m\}$ a set of QC models, such that, for all i , $M_i \subseteq \text{GrdLt}(\mathcal{L}, D)$; The set of all **satisfied formulas** is

$$\text{SF}(\{M_1, \dots, M_n\}) := \{\theta \in \text{Fm}(\mathcal{L}) : \text{for all } i, M_i \models_s \theta\};$$

- The well known Galois connection yields

$$\Delta \subseteq \text{SF}(\text{QC}(\mathcal{L}, \Delta, D));$$

Minimal QC Models

Minimal QC Models

- Let \mathcal{L} be a language and Δ a set of formulas in \mathcal{L} ;

Minimal QC Models

- Let \mathcal{L} be a language and Δ a set of formulas in \mathcal{L} ; The set of **minimal QC models with domain D** is

$$\text{MQC}(\mathcal{L}, \Delta, D) := \{M \in \text{QC}(\mathcal{L}, \Delta, D) : \\ M' \subset M \text{ implies } M' \notin \text{QC}(\mathcal{L}, \Delta, D)\};$$

Minimal QC Models

- Let \mathcal{L} be a language and Δ a set of formulas in \mathcal{L} ; The set of **minimal QC models with domain D** is

$$\text{MQC}(\mathcal{L}, \Delta, D) := \{M \in \text{QC}(\mathcal{L}, \Delta, D) : \\ M' \subset M \text{ implies } M' \notin \text{QC}(\mathcal{L}, \Delta, D)\};$$

- In some sense, these are the models without irrelevant, useless information;

Minimal QC Models

- Let \mathcal{L} be a language and Δ a set of formulas in \mathcal{L} ; The set of **minimal QC models with domain D** is

$$\text{MQC}(\mathcal{L}, \Delta, D) := \{M \in \text{QC}(\mathcal{L}, \Delta, D) : \\ M' \subset M \text{ implies } M' \notin \text{QC}(\mathcal{L}, \Delta, D)\};$$

- In some sense, these are the models without irrelevant, useless information;
- Restricting attention to minimal QC models does not affect reasoning:

Minimal QC Models

- Let \mathcal{L} be a language and Δ a set of formulas in \mathcal{L} ; The set of **minimal QC models with domain D** is

$$\text{MQC}(\mathcal{L}, \Delta, D) := \{M \in \text{QC}(\mathcal{L}, \Delta, D) : \\ M' \subset M \text{ implies } M' \notin \text{QC}(\mathcal{L}, \Delta, D)\};$$

- In some sense, these are the models without irrelevant, useless information;
- Restricting attention to minimal QC models does not affect reasoning:

Minimal Model Theorem

Given a language \mathcal{L} , a set of \mathcal{L} -formulas Δ and a domain D ,

$$\text{SF}(\text{QC}(\mathcal{L}, \Delta, D)) = \text{SF}(\text{MQC}(\mathcal{L}, \Delta, D));$$

Example I

Example I

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;

Example I

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
- Let

$$\Delta = \{\neg P(a) \vee P(b), P(a)\};$$

Example I

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;

- Let

$$\Delta = \{\neg P(a) \vee P(b), P(a)\};$$

- There is only one minimal QC model

$$M = \{P(a), P(b)\};$$

Example I

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;

- Let

$$\Delta = \{\neg P(a) \vee P(b), P(a)\};$$

- There is only one minimal QC model

$$M = \{P(a), P(b)\};$$

- The model $M' = \{P(a), \neg P(a)\}$ is not a QC model of Δ because,

Example I

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;

- Let

$$\Delta = \{\neg P(a) \vee P(b), P(a)\};$$

- There is only one minimal QC model

$$M = \{P(a), P(b)\};$$

- The model $M' = \{P(a), \neg P(a)\}$ is not a QC model of Δ because, if $P(a)$ holds, then $\otimes(\neg P(a) \vee P(b), \neg P(a)) = P(b)$ must also hold;

Example II

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;
- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;
- Let
$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$
- The first sentence says, of course, that “All P 's are Q 's”;

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

$$M_1 = \{P(a), Q(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

$$M_1 = \{P(a), Q(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M_2 = \{P(a), Q(a), Q(b), P(c), \neg P(c), Q(c)\}$$

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;
- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

$$M_1 = \{P(a), Q(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M_2 = \{P(a), Q(a), Q(b), P(c), \neg P(c), Q(c)\}$$

- The following are **not** QC models:

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

$$M_1 = \{P(a), Q(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M_2 = \{P(a), Q(a), Q(b), P(c), \neg P(c), Q(c)\}$$

- The following are **not** QC models:

$$M'_1 = \{P(a), \neg P(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

$$M_1 = \{P(a), Q(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M_2 = \{P(a), Q(a), Q(b), P(c), \neg P(c), Q(c)\}$$

- The following are **not** QC models:

$$M'_1 = \{P(a), \neg P(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M'_2 = \{P(a), \neg P(a), Q(b), P(c), \neg P(c), Q(c)\}$$

Example II

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$\Delta = \{\forall x(\neg P(x) \vee Q(x)), P(a), P(c), \neg P(c)\};$$

- The first sentence says, of course, that “All P 's are Q 's”;
- The following are QC models

$$M_1 = \{P(a), Q(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M_2 = \{P(a), Q(a), Q(b), P(c), \neg P(c), Q(c)\}$$

- The following are **not** QC models:

$$M'_1 = \{P(a), \neg P(a), \neg P(b), P(c), \neg P(c), Q(c)\}$$

$$M'_2 = \{P(a), \neg P(a), Q(b), P(c), \neg P(c), Q(c)\}$$

- Moreover, both M_1 and M_2 are *minimal* QC models;

Example III

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta = \{\forall x \forall y \exists z (\neg P(x, y) \vee Q(z, y)), P(a, b), P(b, b), Q(a, c), \neg Q(a, c)\};$$

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta = \{\forall x \forall y \exists z (\neg P(x, y) \vee Q(z, y)), P(a, b), P(b, b), Q(a, c), \neg Q(a, c)\};$$

- The first sentence imposes the constraint that “Every value of second attribute of P must appear as value of second attribute of Q ”;

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta = \{\forall x \forall y \exists z (\neg P(x, y) \vee Q(z, y)), P(a, b), P(b, b), Q(a, c), \neg Q(a, c)\};$$

- The first sentence imposes the constraint that “Every value of second attribute of P must appear as value of second attribute of Q ”;
- Every model must contain $P(a, b), P(b, b), Q(a, c)$ and $\neg Q(a, c)$;

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta = \{ \forall x \forall y \exists z (\neg P(x, y) \vee Q(z, y)), P(a, b), P(b, b), Q(a, c), \neg Q(a, c) \};$$

- The first sentence imposes the constraint that “Every value of second attribute of P must appear as value of second attribute of Q ”;
- Every model must contain $P(a, b), P(b, b), Q(a, c)$ and $\neg Q(a, c)$;
- Moreover, every minimal model must contain exactly one of $Q(a, b), Q(b, b), Q(c, b)$ or $Q(d, b)$;

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta = \{\forall x \forall y \exists z (\neg P(x, y) \vee Q(z, y)), P(a, b), P(b, b), Q(a, c), \neg Q(a, c)\};$$

- The first sentence imposes the constraint that “Every value of second attribute of P must appear as value of second attribute of Q ”;
- Every model must contain $P(a, b), P(b, b), Q(a, c)$ and $\neg Q(a, c)$;
- Moreover, every minimal model must contain exactly one of $Q(a, b), Q(b, b), Q(c, b)$ or $Q(d, b)$;
- An example of a minimal QC model is

Example III

- Consider $\mathcal{L} = \langle \{a, b, c\}, \{P(2), Q(2)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta = \{\forall x \forall y \exists z (\neg P(x, y) \vee Q(z, y)), P(a, b), P(b, b), Q(a, c), \neg Q(a, c)\};$$

- The first sentence imposes the constraint that “Every value of second attribute of P must appear as value of second attribute of Q ”;
- Every model must contain $P(a, b), P(b, b), Q(a, c)$ and $\neg Q(a, c)$;
- Moreover, every minimal model must contain exactly one of $Q(a, b), Q(b, b), Q(c, b)$ or $Q(d, b)$;
- An example of a minimal QC model is

$$M_1 = \{P(a, b), P(b, b), Q(a, c), \neg Q(a, c), \\ Q(a, b), Q(d, a), Q(b, c), Q(c, d)\};$$

Example IV

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;
- Every minimal model must contain

{

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;
- Every minimal model must contain

$$\{P(a, b, c), \neg P(a, b, c), P(a, d, e),$$

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;
- Every minimal model must contain

$$\{P(a, b, c), \neg P(a, b, c), P(a, d, e), P(a, d, c), P(a, b, e),$$

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;
- Every minimal model must contain

$$\{P(a, b, c), \neg P(a, b, c), P(a, d, e), P(a, d, c), P(a, b, e),$$

- $P(a, d, c)$ and $P(a, b, e)$ are in each model because of the multi-valued dependency;

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;
- Every minimal model must contain

$$\{ P(a, b, c), \neg P(a, b, c), P(a, d, e), P(a, d, c), P(a, b, e), \\ \neg P(a, a, c), \neg P(a, c, c), \neg P(a, d, c), \neg P(a, e, c) \}$$

- $P(a, d, c)$ and $P(a, b, e)$ are in each model because of the multi-valued dependency;

Example IV

- Consider $\mathcal{L} = \langle \{a, b, c, d, e\}, \{P(3)\} \rangle$ and $D = \{a, b, c, d, e\}$;
- Let

$$\Delta = \{ \forall x \forall y \forall z \forall u \forall v (\neg P(x, y, z) \vee \neg P(x, u, v) \vee P(x, y, v)), \\ P(a, b, c), \neg P(a, b, c), P(a, d, e) \};$$

- The first sentence presents a multi-valued dependency of the second and third attributes of P on the first attribute;
- Every minimal model must contain

$$\{ P(a, b, c), \neg P(a, b, c), P(a, d, e), P(a, d, c), P(a, b, e), \\ \neg P(a, a, c), \neg P(a, c, c), \neg P(a, d, c), \neg P(a, e, c) \}$$

- $P(a, d, c)$ and $P(a, b, e)$ are in each model because of the multi-valued dependency;
- The negated atoms are also forced into the model because of the same dependency;

Conflict Base and Model Inconsistency

Conflict Base and Model Inconsistency

- We now introduce a measure for the inconsistency present in a knowledge base;

Conflict Base and Model Inconsistency

- We now introduce a measure for the inconsistency present in a knowledge base;
- Depending on the language \mathcal{L} used and the domain D , a knowledge base may have many different minimal QC models;

Conflict Base and Model Inconsistency

- We now introduce a measure for the inconsistency present in a knowledge base;
- Depending on the language \mathcal{L} used and the domain D , a knowledge base may have many different minimal QC models;
- The measure will be a ratio between 0 and 1, with denominator the total possible number of inconsistencies in the bistructure;

Conflict Base and Model Inconsistency

- We now introduce a measure for the inconsistency present in a knowledge base;
- Depending on the language \mathcal{L} used and the domain D , a knowledge base may have many different minimal QC models;
- The measure will be a ratio between 0 and 1, with denominator the total possible number of inconsistencies in the bistructure;
- Given a QC model M , its **conflict base** is defined by

$$\text{Cnfl}(M) := \{\alpha : \alpha \in M \text{ and } \neg\alpha \in M\};$$

Conflict Base and Model Inconsistency

- We now introduce a measure for the inconsistency present in a knowledge base;
- Depending on the language \mathcal{L} used and the domain D , a knowledge base may have many different minimal QC models;
- The measure will be a ratio between 0 and 1, with denominator the total possible number of inconsistencies in the bistructure;
- Given a QC model M , its **conflict base** is defined by

$$\text{Cnfl}(M) := \{\alpha : \alpha \in M \text{ and } \neg\alpha \in M\};$$

- Recalling the finiteness hypothesis, we define the **measure of inconsistency** for a model M in the context of \mathcal{L} and D (i.e., $\text{Cnfl}(M) \subseteq \text{GrdAt}(\mathcal{L}, D)$) by

$$\text{ModInc}(M, \mathcal{L}, D) := \frac{|\text{Cnfl}(M)|}{|\text{GrdAt}(\mathcal{L}, D)|};$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;
- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;
- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

- We have

$$|\text{GrdAt}(\mathcal{L}, D)| =$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

- We have

$$|\text{GrdAt}(\mathcal{L}, D)| = 12;$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

- We have

$$|\text{GrdAt}(\mathcal{L}, D)| = 12;$$

- Moreover,

$$\text{Cnfl}(M) =$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

- We have

$$|\text{GrdAt}(\mathcal{L}, D)| = 12;$$

- Moreover,

$$\text{Cnfl}(M) = \{P(a, a)\};$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

- We have

$$|\text{GrdAt}(\mathcal{L}, D)| = 12;$$

- Moreover,

$$\text{Cnfl}(M) = \{P(a, a)\};$$

- Therefore,

$$\text{ModInc}(M, \mathcal{L}, D) =$$

Example of Calculating $\text{ModInc}(M, \mathcal{L}, D)$

- Consider $\mathcal{L} = \langle \{\}, \{P(2), R(1)\} \rangle$ and $D = \{a, b, c\}$;

- Let

$$M = \{P(a, a), \neg P(a, a), R(a), \neg R(b), P(b, c)\};$$

- We have

$$|\text{GrdAt}(\mathcal{L}, D)| = 12;$$

- Moreover,

$$\text{Cnfl}(M) = \{P(a, a)\};$$

- Therefore,

$$\text{ModInc}(M, \mathcal{L}, D) = \frac{1}{12};$$

Basic Properties of the Measure

Basic Properties of the Measure

Anti-monotonicity Laws

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 =$$

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\}$$

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 =$$

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$
- Therefore,

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$
 - Therefore, $\text{ModInc}(M_1, \mathcal{L}, D)$

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$
 - Therefore, $\text{ModInc}(M_1, \mathcal{L}, D) > \text{ModInc}(M_2, \mathcal{L}, D)$;

Basic Properties of the Measure

Anti-monotonicity Laws

- If $\mathcal{L}_1 \subseteq \mathcal{L}_2$, then $\text{ModInc}(M, \mathcal{L}_2, D) \leq \text{ModInc}(M, \mathcal{L}_1, D)$;
- If $D_1 \subseteq D_2$, then $\text{ModInc}(M, \mathcal{L}, D_2) \leq \text{ModInc}(M, \mathcal{L}, D_1)$;
- For fixed \mathcal{L} and D , if Δ contains only ground formulas and $M_1, M_2 \in \text{MQC}(\mathcal{L}, \Delta, D)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D);$$

- The property outlined in the previous bullet **does not hold** in general; The following is a counterexample:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$
 - Therefore, $\text{ModInc}(M_1, \mathcal{L}, D) > \text{ModInc}(M_2, \mathcal{L}, D)$;

Preferred QC Models

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ;

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ; Let Δ be a set of \mathcal{L} -formulas;

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ; Let Δ be a set of \mathcal{L} -formulas; Define the set $\text{PQC}(\mathcal{L}, \Delta, D)$ of **preferred QC models of Δ with domain D** by

$$\text{PQC}(\mathcal{L}, \Delta, D) = \{M \in \text{MQC}(\mathcal{L}, \Delta, D) : \\ \text{for all } M' \in \text{MQC}(\mathcal{L}, \Delta, D), |\text{Cnfl}(M)| \leq |\text{Cnfl}(M')|\};$$

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ; Let Δ be a set of \mathcal{L} -formulas; Define the set $\text{PQC}(\mathcal{L}, \Delta, D)$ of **preferred QC models of Δ with domain D** by

$$\text{PQC}(\mathcal{L}, \Delta, D) = \{M \in \text{MQC}(\mathcal{L}, \Delta, D) : \\ \text{for all } M' \in \text{MQC}(\mathcal{L}, \Delta, D), |\text{Cnfl}(M)| \leq |\text{Cnfl}(M')|\};$$

- Revisiting **the preceding example**, recall

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ; Let Δ be a set of \mathcal{L} -formulas; Define the set $\text{PQC}(\mathcal{L}, \Delta, D)$ of **preferred QC models of Δ with domain D** by

$$\text{PQC}(\mathcal{L}, \Delta, D) = \{M \in \text{MQC}(\mathcal{L}, \Delta, D) : \text{for all } M' \in \text{MQC}(\mathcal{L}, \Delta, D), |\text{Cnfl}(M)| \leq |\text{Cnfl}(M')|\};$$

- Revisiting **the preceding example**, recall
 - $\mathcal{L} = \{\{a\}, \{P(1), Q(1)\}\}; D = \{a, b\};$
 $\Delta = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\};$

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ; Let Δ be a set of \mathcal{L} -formulas; Define the set $\text{PQC}(\mathcal{L}, \Delta, D)$ of **preferred QC models of Δ with domain D** by

$$\text{PQC}(\mathcal{L}, \Delta, D) = \{M \in \text{MQC}(\mathcal{L}, \Delta, D) : \\ \text{for all } M' \in \text{MQC}(\mathcal{L}, \Delta, D), |\text{Cnfl}(M)| \leq |\text{Cnfl}(M')|\};$$

- Revisiting **the preceding example**, recall
 - $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$; $D = \{a, b\}$;
 - $\Delta = \langle \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \rangle$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$

Preferred QC Models

- To focus attention to the “least inconsistent” models we restrict the class of minimal QC models to that of preferred QC models, i.e., those minimal ones with a minimal conflict base:
- Consider a language \mathcal{L} and a domain D ; Let Δ be a set of \mathcal{L} -formulas; Define the set $\text{PQC}(\mathcal{L}, \Delta, D)$ of **preferred QC models of Δ with domain D** by

$$\text{PQC}(\mathcal{L}, \Delta, D) = \{M \in \text{MQC}(\mathcal{L}, \Delta, D) : \text{for all } M' \in \text{MQC}(\mathcal{L}, \Delta, D), |\text{Cnfl}(M)| \leq |\text{Cnfl}(M')|\};$$

- Revisiting **the preceding example**, recall
 - $\mathcal{L} = \langle \{a\}, \{P(1), Q(1)\} \rangle$; $D = \{a, b\}$;
 - $\Delta = \{\exists x P(x), \neg P(a) \vee Q(a), \neg Q(a)\}$;
 - Then $\text{MQC}(\mathcal{L}, \Delta, D) = \{M_1, M_2\}$, with

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\} \text{ and } M_2 = \{P(b), \neg P(a), \neg Q(a)\};$$
 - Therefore, $\text{PQC}(\mathcal{L}, \Delta, D) = \{M_2\}$;

Minimal vs. of Minimal Conflict Base

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base:

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;
 - If $M = \{P(a), P(b)\}$, then $M \in \text{QC}(\mathcal{L}, \Delta, D)$, but, clearly, $M \notin \text{MQC}(\mathcal{L}, \Delta, D)$;

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;
 - If $M = \{P(a), P(b)\}$, then $M \in \text{QC}(\mathcal{L}, \Delta, D)$, but, clearly, $M \notin \text{MQC}(\mathcal{L}, \Delta, D)$;
 - However, $|\text{Cnfl}(M)| = 0$, thus, M is of minimal conflict base;

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;
 - If $M = \{P(a), P(b)\}$, then $M \in \text{QC}(\mathcal{L}, \Delta, D)$, but, clearly, $M \notin \text{MQC}(\mathcal{L}, \Delta, D)$;
 - However, $|\text{Cnfl}(M)| = 0$, thus, M is of minimal conflict base;
- We focused attention on $\text{MQC}(\mathcal{L}, \Delta, D)$ to avoid inclusion of superfluous information;

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;
 - If $M = \{P(a), P(b)\}$, then $M \in \text{QC}(\mathcal{L}, \Delta, D)$, but, clearly, $M \notin \text{MQC}(\mathcal{L}, \Delta, D)$;
 - However, $|\text{Cnfl}(M)| = 0$, thus, M is of minimal conflict base;
- We focused attention on $\text{MQC}(\mathcal{L}, \Delta, D)$ to avoid inclusion of superfluous information; Recall, also, that we do not lose (from the deductive point of view) any information;

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;
 - If $M = \{P(a), P(b)\}$, then $M \in \text{QC}(\mathcal{L}, \Delta, D)$, but, clearly, $M \notin \text{MQC}(\mathcal{L}, \Delta, D)$;
 - However, $|\text{Cnfl}(M)| = 0$, thus, M is of minimal conflict base;
- We focused attention on $\text{MQC}(\mathcal{L}, \Delta, D)$ to avoid inclusion of superfluous information; Recall, also, that we do not lose (from the deductive point of view) any information;
- For a particular class of formulas, all minimal models are preferred:

Minimal vs. of Minimal Conflict Base

- There exist non-minimal QC models which possess a minimal conflict base: The following is an example:
 - Consider $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $D = \{a, b\}$;
 - Let $\Delta = \{P(a)\}$;
 - If $M = \{P(a), P(b)\}$, then $M \in \text{QC}(\mathcal{L}, \Delta, D)$, but, clearly, $M \notin \text{MQC}(\mathcal{L}, \Delta, D)$;
 - However, $|\text{Cnfl}(M)| = 0$, thus, M is of minimal conflict base;
- We focused attention on $\text{MQC}(\mathcal{L}, \Delta, D)$ to avoid inclusion of superfluous information; Recall, also, that we do not lose (from the deductive point of view) any information;
- For a particular class of formulas, all minimal models are preferred:

Theorem (Minimal Models for Universal Formulas)

Given \mathcal{L} , Δ and D , if for all $\theta \in \Delta$, θ does not have any existential quantifiers, then $\text{PQC}(\mathcal{L}, \Delta, D) = \text{MQC}(\mathcal{L}, \Delta, D)$.

Non-Standard Behavior of Existential Quantification

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

$$\Delta_2 = \{ P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a) \};$$

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

$$\Delta_2 = \{ P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a) \};$$

- In classical logic, since $D = \{a, b\}$, the sentences $\exists x P(x)$ and $P(a) \vee P(b)$ would be interchangeable;

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

$$\Delta_2 = \{ P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a) \};$$

- In classical logic, since $D = \{a, b\}$, the sentences $\exists x P(x)$ and $P(a) \vee P(b)$ would be interchangeable;
- But in this case, setting

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{\exists xP(x), \neg P(a) \vee Q(a), \neg Q(a)\};$$

$$\Delta_2 = \{P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a)\};$$

- In classical logic, since $D = \{a, b\}$, the sentences $\exists xP(x)$ and $P(a) \vee P(b)$ would be interchangeable;
- But in this case, setting

$$M_1 = \{P(a), \neg P(a), Q(a), \neg Q(a)\};$$

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

$$\Delta_2 = \{ P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a) \};$$

- In classical logic, since $D = \{a, b\}$, the sentences $\exists x P(x)$ and $P(a) \vee P(b)$ would be interchangeable;
- But in this case, setting

$$M_1 = \{ P(a), \neg P(a), Q(a), \neg Q(a) \};$$

$$M_2 = \{ P(b), \neg P(a), \neg Q(a) \};$$

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

$$\Delta_2 = \{ P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a) \};$$

- In classical logic, since $D = \{a, b\}$, the sentences $\exists x P(x)$ and $P(a) \vee P(b)$ would be interchangeable;
- But in this case, setting

$$M_1 = \{ P(a), \neg P(a), Q(a), \neg Q(a) \};$$

$$M_2 = \{ P(b), \neg P(a), \neg Q(a) \};$$

we get $\text{MQC}(\mathcal{L}, \Delta_1, D) = \{M_1, M_2\}$, whereas
 $\text{MQC}(\mathcal{L}, \Delta_2, D) = \{M_2\}$;

Non-Standard Behavior of Existential Quantification

- Consider $\mathcal{L} = \langle \{a, b\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b\}$;
- Consider, also,

$$\Delta_1 = \{ \exists x P(x), \neg P(a) \vee Q(a), \neg Q(a) \};$$

$$\Delta_2 = \{ P(a) \vee P(b), \neg P(a) \vee Q(a), \neg Q(a) \};$$

- In classical logic, since $D = \{a, b\}$, the sentences $\exists x P(x)$ and $P(a) \vee P(b)$ would be interchangeable;
- But in this case, setting

$$M_1 = \{ P(a), \neg P(a), Q(a), \neg Q(a) \};$$

$$M_2 = \{ P(b), \neg P(a), \neg Q(a) \};$$

we get $\text{MQC}(\mathcal{L}, \Delta_1, D) = \{M_1, M_2\}$, whereas

$$\text{MQC}(\mathcal{L}, \Delta_2, D) = \{M_2\};$$

- The discrepancy is of course due to the difference between the **semantics of disjunction** as compared to that of classical logic;

Uniformity of Inconsistency Measure

Uniformity of Inconsistency Measure

- For fixed \mathcal{L} and D ,

Uniformity of Inconsistency Measure

- For fixed \mathcal{L} and D , the inconsistency measure ModInc measures the inconsistency of a set Δ in a *uniform manner*:

Uniformity of Inconsistency Measure

- For fixed \mathcal{L} and D , the inconsistency measure ModInc measures the inconsistency of a set Δ in a *uniform manner*: If $M_1, M_2 \in \text{PQC}(\mathcal{L}, \Delta, D)$, then

Uniformity of Inconsistency Measure

- For fixed \mathcal{L} and D , the inconsistency measure ModInc measures the inconsistency of a set Δ in a *uniform manner*: If $M_1, M_2 \in \text{PQC}(\mathcal{L}, \Delta, D)$, then $|\text{Cnfl}(M_1)| = |\text{Cnfl}(M_2)|$, whence

Uniformity of Inconsistency Measure

- For fixed \mathcal{L} and D , the inconsistency measure ModInc measures the inconsistency of a set Δ in a *uniform manner*: If $M_1, M_2 \in \text{PQC}(\mathcal{L}, \Delta, D)$, then $|\text{Cnfl}(M_1)| = |\text{Cnfl}(M_2)|$, whence $\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D)$;

Uniformity of Inconsistency Measure

- For fixed \mathcal{L} and D , the inconsistency measure ModInc measures the inconsistency of a set Δ in a *uniform manner*: If $M_1, M_2 \in \text{PQC}(\mathcal{L}, \Delta, D)$, then $|\text{Cnfl}(M_1)| = |\text{Cnfl}(M_2)|$, whence $\text{ModInc}(M_1, \mathcal{L}, D) = \text{ModInc}(M_2, \mathcal{L}, D)$;
- Thus, even if there exist many different preferred QC models of Δ , all such models have identical inconsistency measures;

An Example

An Example

- Example:

An Example

- Example:

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;

An Example

- Example:

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

An Example

- **Example:**

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\Delta_1 = \{P(a) \wedge \neg P(a), Q(a) \wedge (Q(b) \vee Q(c) \vee Q(d))\};$$

An Example

- Example:

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\begin{aligned}\Delta_1 &= \{P(a) \wedge \neg P(a), Q(a) \wedge (Q(b) \vee Q(c) \vee Q(d))\}; \\ \Delta_2 &= \{P(a) \wedge \neg P(a), Q(a) \vee Q(b)\};\end{aligned}$$

An Example

- Example:

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\begin{aligned}\Delta_1 &= \{P(a) \wedge \neg P(a), Q(a) \wedge (Q(b) \vee Q(c) \vee Q(d))\}; \\ \Delta_2 &= \{P(a) \wedge \neg P(a), Q(a) \vee Q(b)\};\end{aligned}$$

- We have

An Example

- Example:

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\begin{aligned}\Delta_1 &= \{P(a) \wedge \neg P(a), Q(a) \wedge (Q(b) \vee Q(c) \vee Q(d))\}; \\ \Delta_2 &= \{P(a) \wedge \neg P(a), Q(a) \vee Q(b)\};\end{aligned}$$

- We have

$$\begin{aligned}\text{PQC}(\mathcal{L}, \Delta_1, D) &= \{\{P(a), \neg P(a), Q(a), Q(b)\}, \\ &\quad \{P(a), \neg P(a), Q(a), Q(c)\}, \{P(a), \neg P(a), Q(a), Q(d)\}\};\end{aligned}$$

An Example

- **Example:**

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\begin{aligned}\Delta_1 &= \{P(a) \wedge \neg P(a), Q(a) \wedge (Q(b) \vee Q(c) \vee Q(d))\}; \\ \Delta_2 &= \{P(a) \wedge \neg P(a), Q(a) \vee Q(b)\};\end{aligned}$$

- We have

$$\begin{aligned}PQC(\mathcal{L}, \Delta_1, D) &= \{\{P(a), \neg P(a), Q(a), Q(b)\}, \\ &\quad \{P(a), \neg P(a), Q(a), Q(c)\}, \{P(a), \neg P(a), Q(a), Q(d)\}\}; \\ PQC(\mathcal{L}, \Delta_2, D) &= \{\{P(a), \neg P(a), Q(a)\}, \{P(a), \neg P(a), Q(b)\}\};\end{aligned}$$

An Example

- **Example:**

- Consider $\mathcal{L} = \langle \{a, b, c, d\}, \{P(1), Q(1)\} \rangle$ and $D = \{a, b, c, d\}$;
- Let

$$\begin{aligned}\Delta_1 &= \{P(a) \wedge \neg P(a), Q(a) \wedge (Q(b) \vee Q(c) \vee Q(d))\}; \\ \Delta_2 &= \{P(a) \wedge \neg P(a), Q(a) \vee Q(b)\};\end{aligned}$$

- We have

$$\begin{aligned}\text{PQC}(\mathcal{L}, \Delta_1, D) &= \{\{P(a), \neg P(a), Q(a), Q(b)\}, \\ &\quad \{P(a), \neg P(a), Q(a), Q(c)\}, \{P(a), \neg P(a), Q(a), Q(d)\}\}; \\ \text{PQC}(\mathcal{L}, \Delta_2, D) &= \{\{P(a), \neg P(a), Q(a)\}, \{P(a), \neg P(a), Q(b)\}\};\end{aligned}$$

- Thus, for all $M \in \text{PQC}(\mathcal{L}, \Delta_1, D)$, $\text{ModInc}(M, \mathcal{L}, D) = \frac{1}{8}$ and, for all $M \in \text{PQC}(\mathcal{L}, \Delta_2, D)$, $\text{ModInc}(M, \mathcal{L}, D) = \frac{1}{8}$;

Extrinsic Inconsistency of a Theory

Extrinsic Inconsistency of a Theory

Theorem

Given \mathcal{L} and Δ , if D_1 and D_2 are two domains of the same size and $M_1 \in \text{PQC}(\mathcal{L}, \Delta, D_1)$, $M_2 \in \text{PQC}(\mathcal{L}, \Delta, D_2)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D_1) = \text{ModInc}(M_2, \mathcal{L}, D_2);$$

Extrinsic Inconsistency of a Theory

Theorem

Given \mathcal{L} and Δ , if D_1 and D_2 are two domains of the same size and $M_1 \in \text{PQC}(\mathcal{L}, \Delta, D_1)$, $M_2 \in \text{PQC}(\mathcal{L}, \Delta, D_2)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D_1) = \text{ModInc}(M_2, \mathcal{L}, D_2);$$

- The **extrinsic inconsistency** $\text{ThInc}(\Delta, \mathcal{L})$ of a theory Δ in a language \mathcal{L} is a sequence

$$\langle r_1, r_2, r_3, \dots \rangle,$$

where, for all $n \geq 1$, $r_n = \text{ModInc}(M, \mathcal{L}, D_n)$, where D_n is a domain of size n and $M \in \text{PQC}(\mathcal{L}, \Delta, D_n)$, if such a model exists, and $r_n = *$ (null value), otherwise;

Extrinsic Inconsistency of a Theory

Theorem

Given \mathcal{L} and Δ , if D_1 and D_2 are two domains of the same size and $M_1 \in \text{PQC}(\mathcal{L}, \Delta, D_1)$, $M_2 \in \text{PQC}(\mathcal{L}, \Delta, D_2)$, then

$$\text{ModInc}(M_1, \mathcal{L}, D_1) = \text{ModInc}(M_2, \mathcal{L}, D_2);$$

- The **extrinsic inconsistency** $\text{ThInc}(\Delta, \mathcal{L})$ of a theory Δ in a language \mathcal{L} is a sequence

$$\langle r_1, r_2, r_3, \dots \rangle,$$

where, for all $n \geq 1$, $r_n = \text{ModInc}(M, \mathcal{L}, D_n)$, where D_n is a domain of size n and $M \in \text{PQC}(\mathcal{L}, \Delta, D_n)$, if such a model exists, and $r_n = *$ (null value), otherwise;

- The intention is to measure how the inconsistency of Δ in \mathcal{L} evolves as the size of the domain increases;

Some Examples

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) =$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) =$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) =$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$ and $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$ and $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$

Then $\text{ThInc}(\Delta_1, \mathcal{L}) =$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$ and $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$

$$\text{Then } \text{ThInc}(\Delta_1, \mathcal{L}) = \langle \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \dots \rangle$$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$ and $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$ and $\Delta_2 = \{\exists x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$;
Then $\text{ThInc}(\Delta_1, \mathcal{L}) = \langle \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \dots \rangle$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$ and $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$ and $\Delta_2 = \{\exists x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$;

Then $\text{ThInc}(\Delta_1, \mathcal{L}) = \langle \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \dots \rangle$ and $\text{ThInc}(\Delta_2, \mathcal{L}) =$

Some Examples

- If $\mathcal{L} = \langle \{a\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, 1, \frac{2}{3}, \frac{2}{4}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{a, b, c\}, \{P(1)\} \rangle$ and $\Delta = \{P(a), \neg P(a), P(b), \neg P(b), P(c), \neg P(c)\}$, then

$$\text{ThInc}(\Delta, \mathcal{L}) = \langle *, *, 1, \frac{3}{4}, \frac{3}{5}, \dots \rangle;$$

- If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$ and $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$ and $\Delta_2 = \{\exists x(P(x) \wedge \neg P(x)), \forall x Q(x)\}$;

Then $\text{ThInc}(\Delta_1, \mathcal{L}) = \langle \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \dots \rangle$ and $\text{ThInc}(\Delta_2, \mathcal{L}) = \langle \frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \dots \rangle$;

Some Observations

Some Observations

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If $|\mathcal{C}| = k$, $k \geq 1$, then, for all $1 \leq i < k$, $x_i = *$ and, for all $i \geq k$, $x_i \neq *$.

Some Observations

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If $|\mathcal{C}| = k$, $k \geq 1$, then, for all $1 \leq i < k$, $x_i = *$ and, for all $i \geq k$, $x_i \neq *$.

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If there is an $r_i \in \{r_1, r_2, \dots\}$, such that $r_i = 0$, then $\langle r_1, r_2, \dots \rangle$ is of the form $\langle *, \dots, *, 0, \dots, 0 \rangle$.

Some Observations

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If $|\mathcal{C}| = k$, $k \geq 1$, then, for all $1 \leq i < k$, $x_i = *$ and, for all $i \geq k$, $x_i \neq *$.

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If there is an $r_i \in \{r_1, r_2, \dots\}$, such that $r_i = 0$, then $\langle r_1, r_2, \dots \rangle$ is of the form $\langle *, \dots, *, 0, \dots, 0 \rangle$.

- If $\text{ThInc}(\Delta_1, \mathcal{L}_1) = \langle r_1, r_2, \dots \rangle$ and $\text{ThInc}(\Delta_2, \mathcal{L}_2) = \langle s_1, s_2, \dots \rangle$, we define $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$ iff, for all $i \geq 1$, $r_i \leq s_i$ or $r_i = *$ or $s_i = *$.

Some Observations

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If $|\mathcal{C}| = k$, $k \geq 1$, then, for all $1 \leq i < k$, $x_i = *$ and, for all $i \geq k$, $x_i \neq *$.

Proposition

Suppose $\text{ThInc} = \langle x_1, x_2, \dots \rangle$; If there is an $r_i \in \{r_1, r_2, \dots\}$, such that $r_i = 0$, then $\langle r_1, r_2, \dots \rangle$ is of the form $\langle *, \dots, *, 0, \dots, 0 \rangle$.

- If $\text{ThInc}(\Delta_1, \mathcal{L}_1) = \langle r_1, r_2, \dots \rangle$ and $\text{ThInc}(\Delta_2, \mathcal{L}_2) = \langle s_1, s_2, \dots \rangle$, we define $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$ iff, for all $i \geq 1$, $r_i \leq s_i$ or $r_i = *$ or $s_i = *$.
- $\text{ThInc}(\Delta_1, \mathcal{L}_1) \prec \text{ThInc}(\Delta_2, \mathcal{L}_2)$ abbreviates $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$ and $\text{ThInc}(\Delta_2, \mathcal{L}_2) \not\preceq \text{ThInc}(\Delta_1, \mathcal{L}_1)$;

Some Comparisons

Some Comparisons

- In case $\mathcal{L}_1 = \mathcal{L}_2$, we say that Δ_1 **has smaller than or equal inconsistency as** Δ_2 and write $\Delta_1 \leq_{\text{inc}}^{\mathcal{L}} \Delta_2$;

Some Comparisons

- In case $\mathcal{L}_1 = \mathcal{L}_2$, we say that Δ_1 **has smaller than or equal inconsistency as** Δ_2 and write $\Delta_1 \leq_{\text{inc}}^{\mathcal{L}} \Delta_2$;
- For fixed Δ and \mathcal{L} , it is not necessarily the case that $\text{ThInc}(\Delta, \mathcal{L}) = \langle r_1, r_2, \dots \rangle$ is such that, for all i , if $r_i \neq *$, then $r_i \geq r_{i+1}$;

Some Comparisons

- In case $\mathcal{L}_1 = \mathcal{L}_2$, we say that Δ_1 **has smaller than or equal inconsistency as** Δ_2 and write $\Delta_1 \leq_{\text{inc}}^{\mathcal{L}} \Delta_2$;
 - For fixed Δ and \mathcal{L} , it is not necessarily the case that $\text{ThInc}(\Delta, \mathcal{L}) = \langle r_1, r_2, \dots \rangle$ is such that, for all i , if $r_i \neq *$, then $r_i \geq r_{i+1}$;
- Example:** Consider, for instance, $\mathcal{L} = \langle \{\}, \{P(2), Q(1)\} \rangle$ and $\Delta = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$;

Some Comparisons

- In case $\mathcal{L}_1 = \mathcal{L}_2$, we say that Δ_1 **has smaller than or equal inconsistency as** Δ_2 and write $\Delta_1 \leq_{\text{inc}}^{\mathcal{L}} \Delta_2$;
- For fixed Δ and \mathcal{L} , it is not necessarily the case that $\text{ThInc}(\Delta, \mathcal{L}) = \langle r_1, r_2, \dots \rangle$ is such that, for all i , if $r_i \neq *$, then $r_i \geq r_{i+1}$;

Example: Consider, for instance, $\mathcal{L} = \langle \{\}, \{P(2), Q(1)\} \rangle$ and $\Delta = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$; Then $\text{ThInc}(\Delta, \mathcal{L}) = \langle \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots \rangle$;

Some Comparisons

- In case $\mathcal{L}_1 = \mathcal{L}_2$, we say that Δ_1 **has smaller than or equal inconsistency as** Δ_2 and write $\Delta_1 \leq_{\text{inc}}^{\mathcal{L}} \Delta_2$;
 - For fixed Δ and \mathcal{L} , it is not necessarily the case that $\text{ThInc}(\Delta, \mathcal{L}) = \langle r_1, r_2, \dots \rangle$ is such that, for all i , if $r_i \neq *$, then $r_i \geq r_{i+1}$;
- Example:** Consider, for instance, $\mathcal{L} = \langle \{\}, \{P(2), Q(1)\} \rangle$ and $\Delta = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$; Then $\text{ThInc}(\Delta, \mathcal{L}) = \langle \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots \rangle$;

Proposition

If $\Delta_1 \subseteq \Delta_2$, then $\text{ThInc}(\Delta_1, \mathcal{L}) \preceq \text{ThInc}(\Delta_2, \mathcal{L})$.

Some Comparisons

- In case $\mathcal{L}_1 = \mathcal{L}_2$, we say that Δ_1 **has smaller than or equal inconsistency as** Δ_2 and write $\Delta_1 \leq_{\text{inc}}^{\mathcal{L}} \Delta_2$;
- For fixed Δ and \mathcal{L} , it is not necessarily the case that $\text{ThInc}(\Delta, \mathcal{L}) = \langle r_1, r_2, \dots \rangle$ is such that, for all i , if $r_i \neq *$, then $r_i \geq r_{i+1}$;

Example: Consider, for instance, $\mathcal{L} = \langle \{\}, \{P(2), Q(1)\} \rangle$ and $\Delta = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$; Then $\text{ThInc}(\Delta, \mathcal{L}) = \langle \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots \rangle$;

Proposition

If $\Delta_1 \subseteq \Delta_2$, then $\text{ThInc}(\Delta_1, \mathcal{L}) \preceq \text{ThInc}(\Delta_2, \mathcal{L})$.

Proposition

If $\mathcal{L}_1 = \langle \mathcal{C}_1, \mathcal{P}_1 \rangle$ and $\mathcal{L}_2 = \langle \mathcal{C}_2, \mathcal{P}_2 \rangle$, and $\mathcal{L}_1 \subseteq \mathcal{L}_2$ (meaning $\mathcal{C}_1 \subseteq \mathcal{C}_2$ and $\mathcal{P}_1 \subseteq \mathcal{P}_2$), then $\text{ThInc}(\Delta, \mathcal{L}_2) \preceq \text{ThInc}(\Delta, \mathcal{L}_1)$.

More Comparisons

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- Example

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example**
 - For instance, consider

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle \qquad \Delta_1 = \{P(a)\}$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle$$

$$\mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_1 = \{P(a)\}$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle$$

$$\mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_1 = \{P(a)\}$$

$$\Delta_2 = \{P(a), \neg P(a)\}$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_1 = \{P(a)\}$$

$$\mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_2 = \{P(a), \neg P(a)\}$$

- Then

$$\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example and Counterexample:**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_1 = \{P(a)\}$$

$$\mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_2 = \{P(a), \neg P(a)\}$$

- Then

$$\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example and Counterexample:**
 - For instance, consider

$$\mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_1 = \{P(a)\}$$

$$\mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle$$

$$\Delta_2 = \{P(a), \neg P(a)\}$$

$$\mathcal{L}_3 = \langle \{a\}, \{P(1), Q(1)\} \rangle$$

- Then

$$\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example and Counterexample:**
 - For instance, consider

$$\begin{array}{ll}
 \mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle & \Delta_1 = \{P(a)\} \\
 \mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle & \Delta_2 = \{P(a), \neg P(a)\} \\
 \mathcal{L}_3 = \langle \{a\}, \{P(1), Q(1)\} \rangle & \Delta_3 = \{P(a), \neg P(a), Q(a)\};
 \end{array}$$

- Then

$$\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$$

More Comparisons

- If $\Delta_1 \subseteq \Delta_2$ and $\mathcal{L}_1 \subseteq \mathcal{L}_2$, it does not necessarily follow that $\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$;
- **Example and Counterexample:**
 - For instance, consider

$$\begin{array}{ll}
 \mathcal{L}_1 = \langle \{a\}, \{P(1)\} \rangle & \Delta_1 = \{P(a)\} \\
 \mathcal{L}_2 = \langle \{a\}, \{P(1)\} \rangle & \Delta_2 = \{P(a), \neg P(a)\} \\
 \mathcal{L}_3 = \langle \{a\}, \{P(1), Q(1)\} \rangle & \Delta_3 = \{P(a), \neg P(a), Q(a)\};
 \end{array}$$

- Then

$$\text{ThInc}(\Delta_1, \mathcal{L}_1) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2)$$

- However,

$$\text{ThInc}(\Delta_3, \mathcal{L}_3) \preceq \text{ThInc}(\Delta_2, \mathcal{L}_2);$$

Intrinsic Inconsistency of a Theory

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ;

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency** of Δ is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency** of Δ is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$,

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency of Δ** is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\}\rangle$, $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x(Q(x) \wedge \neg Q(x))\}$

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency of Δ** is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\}\rangle$, $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x(Q(x) \wedge \neg Q(x))\}$ and $\Delta_2 = \{\forall x(P(x) \wedge \neg P(x))\}$,

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency of Δ** is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$, $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x(Q(x) \wedge \neg Q(x))\}$ and $\Delta_2 = \{\forall x(P(x) \wedge \neg P(x))\}$, then $\text{ThInc}(\Delta_1) = \text{ThInc}(\Delta_2) =$

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency of Δ** is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$, $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x(Q(x) \wedge \neg Q(x))\}$ and $\Delta_2 = \{\forall x(P(x) \wedge \neg P(x))\}$, then $\text{ThInc}(\Delta_1) = \text{ThInc}(\Delta_2) = \langle 1, 1, \dots \rangle$,

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency of Δ** is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$, $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x(Q(x) \wedge \neg Q(x))\}$ and $\Delta_2 = \{\forall x(P(x) \wedge \neg P(x))\}$, then $\text{ThInc}(\Delta_1) = \text{ThInc}(\Delta_2) = \langle 1, 1, \dots \rangle$, whereas $\text{ThInc}(\Delta_2, \mathcal{L}) \prec \text{ThInc}(\Delta_1, \mathcal{L})$,

Intrinsic Inconsistency of a Theory

- For Δ, Δ' sets of \mathcal{L} -formulas, Δ is **QC equivalent** to Δ' iff for every M , M is a QC model of Δ iff it is a QC model of Δ' ;

Theorem

If Δ_1 and Δ_2 are QC equivalent sets of formulas in the language \mathcal{L} , then

$$\text{ThInc}(\Delta_1, \mathcal{L}) = \text{ThInc}(\Delta_2, \mathcal{L});$$

- For a fixed theory Δ consider the language \mathcal{L} consisting exactly of the predicate and constant symbols appearing in Δ ; The **intrinsic inconsistency of Δ** is defined by

$$\text{ThInc}(\Delta) := \text{ThInc}(\Delta, \mathcal{L});$$

- Example:** If $\mathcal{L} = \langle \{\}, \{P(1), Q(1)\} \rangle$, $\Delta_1 = \{\forall x(P(x) \wedge \neg P(x)), \forall x(Q(x) \wedge \neg Q(x))\}$ and $\Delta_2 = \{\forall x(P(x) \wedge \neg P(x))\}$, then $\text{ThInc}(\Delta_1) = \text{ThInc}(\Delta_2) = \langle 1, 1, \dots \rangle$, whereas $\text{ThInc}(\Delta_2, \mathcal{L}) \prec \text{ThInc}(\Delta_1, \mathcal{L})$, the addition of Q reducing the inconsistency of Δ_2 ;

Comparing Intrinsic Theory Inconsistencies

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;
- Any inconsistent theory has greater inconsistency than any consistent theory;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;
- Any inconsistent theory has greater inconsistency than any consistent theory;
- **Examples:** Let us fix $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;
- Any inconsistent theory has greater inconsistency than any consistent theory;
- **Examples:** Let us fix $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_1 = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has one preferred QC model

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;
- Any inconsistent theory has greater inconsistency than any consistent theory;
- **Examples:** Let us fix $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_1 = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has one preferred QC model $M_1 = \{P(a, a), \neg P(a, a), \dots, P(c, c), \neg P(c, c)\}$;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;
- Any inconsistent theory has greater inconsistency than any consistent theory;
- **Examples:** Let us fix $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_1 = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has one preferred QC model $M_1 = \{P(a, a), \neg P(a, a), \dots, P(c, c), \neg P(c, c)\}$; Therefore, $\text{ModInc}(M_1, \mathcal{L}, D) = \frac{9}{9} = 1$;

Comparing Intrinsic Theory Inconsistencies

- A theory Δ_1 **has smaller than or equal inconsistency as** Δ_2 , written $\Delta_1 \preceq \Delta_2$, iff $\text{ThInc}(\Delta_1) \preceq \text{ThInc}(\Delta_2)$;
- All consistent theories have an intrinsic inconsistency $\langle *, \dots, *, 0, \dots, 0 \rangle$, where the number of *'s depends on the number of constants in the language; Thus, they have the same inconsistency;
- Any inconsistent theory has greater inconsistency than any consistent theory;
- **Examples:** Let us fix $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_1 = \{\forall x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has one preferred QC model $M_1 = \{P(a, a), \neg P(a, a), \dots, P(c, c), \neg P(c, c)\}$; Therefore, $\text{ModInc}(M_1, \mathcal{L}, D) = \frac{9}{9} = 1$; M_1 is totally inconsistent;

Examples (Cont'd)

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$; So, $\text{ModInc}(M_{31}, \mathcal{L}, D) = \frac{3}{9} = \frac{1}{3}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$; So, $\text{ModInc}(M_{31}, \mathcal{L}, D) = \frac{3}{9} = \frac{1}{3}$;
 - The theory $\Delta_4 = \{\exists x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$; So, $\text{ModInc}(M_{31}, \mathcal{L}, D) = \frac{3}{9} = \frac{1}{3}$;
 - The theory $\Delta_4 = \{\exists x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{41} = \{P(b, a), \neg P(b, a), P(b, b), \neg P(b, b), P(b, c), \neg P(b, c)\}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$; So, $\text{ModInc}(M_{31}, \mathcal{L}, D) = \frac{3}{9} = \frac{1}{3}$;
 - The theory $\Delta_4 = \{\exists x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{41} = \{P(b, a), \neg P(b, a), P(b, b), \neg P(b, b), P(b, c), \neg P(b, c)\}$; Therefore, $\text{ModInc}(M_{41}, \mathcal{L}, D) = \frac{1}{3}$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$; So, $\text{ModInc}(M_{31}, \mathcal{L}, D) = \frac{3}{9} = \frac{1}{3}$;
 - The theory $\Delta_4 = \{\exists x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{41} = \{P(b, a), \neg P(b, a), P(b, b), \neg P(b, b), P(b, c), \neg P(b, c)\}$; Therefore, $\text{ModInc}(M_{41}, \mathcal{L}, D) = \frac{1}{3}$;
- For the theories above, the intrinsic complexities are, respectively, $\langle 1, 1, \dots \rangle$, $\langle 1, \frac{1}{4}, \frac{1}{9}, \dots \rangle$, $\langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle$ and $\langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle$;

Examples (Cont'd)

- **Examples:** Still considering $\mathcal{L} = \langle \{\}, \{P(2)\} \rangle$ and $D = \{a, b, c\}$;
 - The theory $\Delta_2 = \{\exists x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One of them is $M_{21} = \{P(a, b), \neg P(a, b)\}$; Therefore, $\text{ModInc}(M_{21}, \mathcal{L}, D) = \frac{1}{9}$;
 - The theory $\Delta_3 = \{\forall x \exists y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{31} = \{P(a, a), \neg P(a, a), P(b, c), \neg P(b, c), P(c, a), \neg P(c, a)\}$; So, $\text{ModInc}(M_{31}, \mathcal{L}, D) = \frac{3}{9} = \frac{1}{3}$;
 - The theory $\Delta_4 = \{\exists x \forall y (P(x, y) \wedge \neg P(x, y))\}$ has 9 preferred QC models; One is $M_{41} = \{P(b, a), \neg P(b, a), P(b, b), \neg P(b, b), P(b, c), \neg P(b, c)\}$; Therefore, $\text{ModInc}(M_{41}, \mathcal{L}, D) = \frac{1}{3}$;
- For the theories above, the intrinsic complexities are, respectively, $\langle 1, 1, \dots \rangle$, $\langle 1, \frac{1}{4}, \frac{1}{9}, \dots \rangle$, $\langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle$ and $\langle 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle$;
- So

$$\text{ThInc}(\Delta_2) <_{\text{inc}} \text{ThInc}(\Delta_3) =_{\text{inc}} \text{ThInc}(\Delta_4) <_{\text{inc}} \text{ThInc}(\Delta_1);$$

A Final Theorem

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \left\langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \right\rangle.$$

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

$$\Delta = \{ \forall x (P_1(x) \wedge \dots \wedge P_s(x)), \neg P_1(a), \dots, \neg P_r(a) \};$$

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

$$\Delta = \{ \forall x (P_1(x) \wedge \dots \wedge P_s(x)), \neg P_1(a), \dots, \neg P_r(a) \};$$

- What is the number $|\text{GrdAt}(\mathcal{L}, D_n)|$?

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

$$\Delta = \{ \forall x (P_1(x) \wedge \dots \wedge P_s(x)), \neg P_1(a), \dots, \neg P_r(a) \};$$

- What is the number $|\text{GrdAt}(\mathcal{L}, D_n)|$? ns ;

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \left\langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \right\rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

$$\Delta = \{ \forall x (P_1(x) \wedge \dots \wedge P_s(x)), \neg P_1(a), \dots, \neg P_r(a) \};$$

- What is the number $|\text{GrdAt}(\mathcal{L}, D_n)|$? ns ;
- What is the number $|\text{Cnfl}(M)|$ if $M \in \text{PQC}(\mathcal{L}, \Delta, D_n)$?

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \left\langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \right\rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

$$\Delta = \{ \forall x (P_1(x) \wedge \dots \wedge P_s(x)), \neg P_1(a), \dots, \neg P_r(a) \};$$

- What is the number $|\text{GrdAt}(\mathcal{L}, D_n)|$? ns ;
- What is the number $|\text{Cnfl}(M)|$ if $M \in \text{PQC}(\mathcal{L}, \Delta, D_n)$? r ;

A Final Theorem

Existence Theorem

For any fraction $\frac{r}{s}$, with $r \leq s$, there exists a theory Δ , such that

$$\text{ThInc}(\Delta) = \langle \frac{r}{s}, \frac{r}{2s}, \frac{r}{3s}, \dots \rangle.$$

- Simply take $\mathcal{L} = \langle \{a\}, \{P_1(1), \dots, P_s(1)\} \rangle$;

$$\Delta = \{ \forall x (P_1(x) \wedge \dots \wedge P_s(x)), \neg P_1(a), \dots, \neg P_r(a) \};$$

- What is the number $|\text{GrdAt}(\mathcal{L}, D_n)|$? ns ;
- What is the number $|\text{Cnfl}(M)|$ if $M \in \text{PQC}(\mathcal{L}, \Delta, D_n)$? r ;
- Thus $r_n = \text{ModInc}(M, \mathcal{L}, D_n) = \frac{r}{ns}$;

Many Thanks!

MANY THANKS for

- the **hospitality**
- your **attention during the seminar!**